

LA-UR-21-24254

Approved for public release; distribution is unlimited.

Title: w19_OMECS Scientific Highlight

Author(s): Zelenay, Piotr

Intended for: Report

Issued: 2021-05-03

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

w19_OMECE Scientific Highlight

❑ Scientific Achievement


- Used DFT to identify likely experimental limitations for realizing high HER activity for DPA-based organic molecular electrocatalysts (OMECE) systems, directly guiding future experiments

❑ Significance and Impact

- OMECEs promise low-cost, durable, systems for hydrogen production and energy applications
- Design of OMECEs structures remain challenging due to lack of fundamental understanding
- Density functional theory (DFT) has been proposed to provide strong insights into HER activity
- DFT has identified non-graphitic/defected C as likely cause for low-activity found experimentally

❑ Research Details

- DFT calculations on the binding energy of HER intermediates for predicting reduction potentials and H adsorption energy as activity descriptors
- LANL Institutional Computing (HPC) Resources essential for successful execution of project for VASP and ADF calculations of HER descriptors



DPA+H on monovacancy (circles) containing graphene basal plane – this shifts H-binding energy descriptor from close to ideal to binding H ~1 eV too strongly.